

---

## Review Article

### Cannabis “in utero”: the fetus as a compulsive consumer

Luiz Antonio Del Ciampo<sup>1</sup>, Ieda Regina Lopes Del Ciampo<sup>2</sup>

<sup>1</sup>Department of Puericulture and Pediatrics – Faculty of Medicine of Ribeirão Preto – University of São Paulo – Brazil

<sup>2</sup>Department of Medicine – Federal University of São Carlos – São Paulo – Brazil

---

#### Abstract:

Cannabis sativa, a plant known for millennia for its hallucinogenic and medicinal effects, contains widely consumed psychoactive substances worldwide, with increasing numbers among adolescents and adults, including pregnant women. Due to its potential adverse effects on users, it is of fundamental importance to know and disseminate the harm caused to humans in order to reduce the consumption of this drug. This article presents a review of the physiological and pharmacological characteristics of psychoactive substances present in cannabis, their actions on the endocannabinoid system and the placenta. It also highlights the main clinical repercussions that can occur with the fetus and, in the long run, with children whose mothers used cannabis during pregnancy.

---

**Key Words:** cannabis – cannabinoids – pregnancy – child health – abuse drugs

---

#### Introduction

Since ancient times human beings have been searching in nature for sources of pleasure and/or relief for the suffering and stress of everyday life, experimenting with the most varied substances that, isolated or combined, can be incorporated into their daily habits. Over time, and adopting social rules, these substances were designated as licit or illicit according to their pharmacological properties and effects on the organism [1].

Among the various illicit substances cannabis products has been used for centuries for spiritual, medicinal and nonmedical purposes, and over the past 60 years recreational or non-medicinal use has spread globally by adolescents and young adults, currently being the most commonly used by pregnant women that believe cannabis is natural and safer than other substances [2-6].

Studies have observed increased prevalence of pregnant women using cannabis in different forms. This increase in consumption is directly linked to the legalization that has been taking place in many countries, the increased availability on this drug, and the lack of accurate information on health risks often overlooked by users, causing the medical and social use of cannabis to become more acceptable in the whole world [7-10].

#### Cannabis, cannabinoids and endocannabinoid system

*Cannabis sativa* (also known as marijuana) is a plant originated in Central Asia and currently growing worldwide that contains more than 500 active herbal compounds and can be consumed inhaled, ingested or applied topically to mucous membranes [11-13]. The most common route of use is smoked because it is the most efficient way to achieve the desired psychoactive effects [14], although the effects depend on the

mode of administration, the dose received, the previous experience, the mood state and the social setting in which it is used [14,15].

Cannabis is the generic term used to denote plant-derived psychoactive substances prepared from *C. sativa*, and consists of approximately 60 plant-derivative cannabinoid compounds denominated phytocannabinoids. In addition to these, there are two other types: the endocannabinoids (naturally produced by the body) and the synthetic cannabinoids (eliciting cannabis effects, generally with a higher potency than endocannabinoids) [16]. The most important psychoactive phytocannabinoid is the delta-9-tetrahydrocannabinol (delta9THC), a partial agonist of the endocannabinoid receptors CBR1 and CBR2, the activation of which generates central and peripheral effects [17-21]. The concentration of delta9THC varies in the different strains of cannabis (6% to 20%), and due to its physicochemical properties (small molecular size and lipophilic nature) has high distribution and transport capacity in the human body, quickly crossing the transplacental membranes [2,22-24]. It is estimated that one-third of delta9THC plasma concentration crosses the fetoplacental barriers exposing the fetuses of cannabis using mothers during the prenatal period. Also, there has been an increase in the potency of cannabis over the past 20 years due to the novel ways of cultivating and production of synthetic substances which enhance the possibility of greater adverse effects among users [25,26].

The effects of delta9THC are due to its action on cannabinoid receptors CBR1 and CBR2 - widely distributed throughout various organs and tissues that, together with endogenously produced cannabinoids and the regulatory enzymes of synthesis and degradation, constitute the endocannabinoid

system (ECS) [19,27]. The ECS is a neuromodulatory system, detected from the earliest embryonic stage and throughout pre- and postnatal development, responsible for several controls related to the regulation of many important function, since its receptors has been found to play a key role in neuronal progenitor cell proliferation, pyramidal specification, axon patterning modulation of dendritic arbor, and promotion of neuronal differentiation [5,20,28,29]. Also, the ECS influences metabolism and physiology of multiple systems with its anabolic action, leading to protein and glycogen synthesis and fat deposition [30]. Moreover, plays a fundamental role in pregnancy outcome and fetal development, and over-stimulation of CB1 in the placenta can impair fetal growth by inhibiting cytotrophoblastic proliferation [5,28,31].

The endocannabinoid receptors can be found on both presynaptic neurons and postsynaptic locations and are densely expressed in various regions within the brain [4], and its key function in mature neurons is modulate the release of neurotransmitters [18]. These receptors has been detected in the fetal human brain as early as 14 weeks of gestation and changes across development indicating important roles in fetal central nervous system (CNS) development events [5,32] such as neuronal proliferation, migration and synaptogenesis [18,33]. Cannabis exposure during pregnancy has the potential to induce supra-physiological stimulation of the ECS, wich can break the ontogeny of endogenous endocannabinoid signalling and compromise the synaptogenesis and the development of neural interconections [17,31,33,34].

Due to the mode of cultivation, preparation and handling some products derived from cannabis can deliver chemical and biological contaminants to the user such as pesticides, yeasts, bacteria, metals and solvents, increasing the harmfulness and the possibility of synergistic effects with other substances [35-38]. Cannabis, with its hundreds of compounds, has the property of interacting with many substances and when consumed in inhaled form, pyrolysis can produce more harmful substances [35].

**Pharmacokinetics of cannabinoids and pregnancy**

Cannabis consumption during pregnancy may negatively impact the fetus since the effects may be subtle and do not be detectable for many months to years after birth [28]. Cannabis can affect the normal transport functions and physiologic status of the placenta where it arrives by passive diffusion or, less commonly, through active transport or pinocytosis [2,19], and its concentrations varies by the permeability and biological capacity of the placenta. In addition, when cannabis is smoked, serum carbon monoxide concentrations in the pregnant woman is very high, resulting potentially in impaired maternal respiratory gas exchange and subsequent adverse effect on the fetus [2].

Other effects observed are increased resistance and pulsatility index of the uterine artery, with resulting potential effects on uterine blood flow, resulting in increased placental resistance and reduced placental circulation [2], besides impair fetal growth by inhibiting cytotrophoblastic proliferation [36].

**Effects on pregnant woman and fetus**

For pregnant women, as for any adult, cannabis use can cause many short-term and long-term biopsychosocial problems (impairment of motor coordination and memory, altered judgement, risk of psychosis, myocardial infaction, arrhythmias, acute coronary syndrome, perpheral arteritis, stroke and cerebrovascular accident) besides predispose to the development of addition and adversely affect a mother’s ability to care for the child [39,40].

Several studies have reported associations between cannabis use during pregnancy and fetal outcomes, congenital anomalies and teratogenic effects [19,34,41]. The effects may vary depending on duration, route administration, timing and overall magnitude of exposure, and the extent to wich the fetus and the fetal SNC are exposed [19], since prenatal exposure to exogenous cannabinoids can modify the maturation of neurotransmitter system and their functions through the activation of receptors that arise early in the developing brain [5].

Alterations on fetal neuroendocrine and neurotransmitters systems, decrease in insulin-like growth factors and effects on direct effects on central nervous system are the main mechanisms that compromise the fetus during pregnancy. The main effects on the fetus caused by cannabis use by pregnant women are presented in Table 1.

**Table 1. Effects of cannabis on the fetus**

Growth restriction
Decreased birth weight
Small for gestatinal age
Decreased head circumference
Reduced grey matter volume
Lower apgar scores
Preterm delivery
More likely to require admission to neonatal intensive care unit
stillbirth
Risk of congenital malformations
Ref. 2,3,8,14,17,31-33,42-56

**Long term effects on the child**

In addition to the immediate effects on the intrauterine environment, the consumption of cannabis use by pregnant women may lead to development abnormalities: tremors, startle, altered responses to visual stimuli, and high-pitched cry after birth [14]. In children have been observed decreased attention and motivation, gaps in problem-solving skill and memory, increased depressive and anxiety symptoms, impulsivity, aggressiveness, disturbed sleep [17,32,46,49,57,58].

**Conclusion**

All early chemical exposure can result in injuries to the fetus that are expressed as changes in postnatal development [18]. External stimuli can affect embryonic signalling systems, resulting in long-lasting or even permanent alterations to

organogenesis leading to the in uteroadaptation of the fetus to an anticipated non-physiological environment after birth [46]. Therefore, in addition to its harmful effects, cannabis can enhance the placental barrier permeability to pharmacologic agents and recreational substances, potentially placing the fetus at risk from other agents or drugs [2].

It is urgent to spread the potential repercussions of cannabis use on maternal and fetal health during pregnancy [44]. There is mounting evidence to suggest that perinatal cannabis exposure can have negative effects and the combination of increasing trends in consume, potency, and access poses an important public health and clinical practice challenge [33,59]. Whereas the general population increasingly views cannabis as safe, measures to disseminate reliable information on the harm caused by cannabis use during pregnancy are urgently needed, preventing thousands of children from being exposed to the harmful effects that begin during the intrauterine period [18,53,61].

Preventive programs should provide education about the potential adverse effects on the health of pregnant women and their future or current fetus or newborn [62,63]. During the initial prenatal all women should be asked about use of cannabis and medical community should advise pregnant women to avoid perinatal cannabis exposure [36,64]. Besides that it is very important education of health providers in the care and management of women with evidence of drug use before, during and after pregnancy.

## References

1. Pisanti S, Bifulco M. Medical cannabis: a plurimillennial history of an evergreen. *J Cell Physiol* 2019;234:8342-8351.
2. Ryan AS, Ammerman SD, FSAHM, O'Connor ME. Marijuana use during pregnancy and breastfeeding: implications for neonatal and childhood outcomes. *Pediatrics* 2018;142:1-17.
3. El Marroun H, Brown QL, Lund IO, Coleman-Cowger VH, Loree AM, Chawla D et al. An epidemiological, developmental and clinical overview of cannabis during pregnancy. *Prev Med* 2018;116:1-5.
4. Cohen K, Weizamn A, Weinstein A. Positive and negative effects of cannabis and cannabinoids in health. *Clin Pharmacol Ther* 2019;105:1139-1147.
5. Velez ML, Jordan CJ, Jansson LM. (2019) Cannabis use disorder during the perinatal period. In: Montoya I., Weiss S. (eds) *Cannabis Use Disorders*. Springer Nature Switzerland.
6. Volkow ND, Han B, Compton WM. Marijuana use during stages of pregnancy in the United States. *Ann Intern Med* 2017;166:763-764.
7. Chabarria KC, Racusin DA, Antony KM, Kahr M, Suter MA, Mastrobattista JM et al. Marijuana use and its effects in pregnancy. *Am J Obstet Gynecol* 2016;215:506.e1-506.e7.
8. Gunn JKL, Rosales CB, Center KE, Nuñez A, Gibson SJ, Christ C et al. Prenatal exposure to cannabis and maternal and child health outcomes: a systematic review and meta-analysis. *BMJ Open* 2016;6:1-10.
9. Merlo P, Stahl B, Klinger P. Does cannabis use by the pregnant mother affect the fetus and newborn? *Ped Endocrinol Rev* 2017;15:4-7.
10. Bayrampour H, Zahradnik M, Lisonkova S, Janssen P. Women's perspectives about cannabis use during pregnancy and the postpartum period: an integrative review. *Prev Med* 2019;119:17-23.
11. Peters J, Chien J. Contemporary routes of cannabis consumption: a primer for clinicians. *J Am Osteopath Assoc* 2018;118:67-70.
12. Chadi N, Levy S. What every pediatric gynecologist should know about marijuana use in adolescents. *J Pediatr Adolesc Gynecol* 2019;32:1-5.
13. Qian Y, Gurley BJ, Markowitz JS. The potential for pharmacokinetic interactions between cannabis products and conventional medications. *J Clin Psychopharmacol* 2019;39:462-471.
14. Hall W, Degenhardt L. Adverse health effects of non-medical cannabis use. *Lancet* 2009;374:1383-1390.
15. Strecka JM, Hughesa JR, Klemperer EM, Howard AB, Budney AJ. Modes of cannabis use: A secondary analysis of an intensive longitudinal natural history study. *Addict Behav* 2019;98:1-7.
16. Kim J, Castro A, Lendoiro E, Cruz-Landeira A, López-Rivadulla M, Concheiro M. Detection of in utero cannabis exposure by umbilical cord analysis. *Drug Test Anal* 2018;10:636-643.
17. Wu C, Jew CP, Lu H. Lasting impacts of prenatal cannabis exposure and the role of endogenous cannabinoids in the developing brain. *Future Neurol* 2011;6:459-480.
18. Grant KS, Petroff R, Isoherranen N, Stella N, Burbacher TM. Cannabis use during pregnancy: pharmacokinetics and effects on child development. *Pharmacol Ther* 2018;182:133-151.
19. Thompson R, DeJong K, Lo J. Marijuana use in pregnancy: a review. *Obstet Gynecol Surv* 2019;74:515-422.
20. Alexandre J, Carmo H, Carvalho F, Silva JP. Synthetic cannabinoids and their impact on neurodevelopmental processes. *Addiction Biology* 2019:e12824. [Epub ahead of print].
21. Carlier J, Huestis MA, Zaami S, Pichini S, Busardò FP. Monitoring the effects of perinatal cannabis and synthetic cannabinoid exposure. *Ther Drug Monit* 2019;jun 28. [Epub ahead of print].
22. Silva JP, Yonamine M. Bioanalytical and methodological challenges in the evaluation of fetal cannabis exposure. *Bioanalysis* 2018;10:713-716.
23. Dinis-Oliveira RJ. Metabolomics of delta9-tetrahydrocannabinol: implications in toxicity. *Drug Metab Rev* 2016;48:80-87.
24. De Luca MA, Di Chiara G, Cadoni C, Lecca D, Orsolini L, Papanti D et al. Cannabis; epidemiological, neurobiological and psychopathological issues: an update. *CNS Neurol Disord Drug Targets* 2017;16:598-609.

25. Alipour A, Patel PB, Shabbir Z, Gabrielson S. Review of the many faces of synthetic cannabinoid toxicities. *Ment Health Clin* 2019;9:93-99.
26. Turner SE, Williams CM, Iversen L. Molecular pharmacology of phytocannabinoids. *Prog Chem Org Nat Prod* 2017;103:61–101.
27. Bonini SA, Premoli M, Tambaro S. *Cannabis sativa*: a comprehensive ethnopharmacological review of a medicinal plant with a long history. *J Ethnopharmacol*. 2018;227:300–315.
28. Maia J, Midão L, Cunha SC, Almada M, Fonseca BM, Braga J et al. Effects of cannabis tetrahydrocannabinol on endocannabinoid homeostasis in human placenta. *Arch Toxicol* 2019;93:649-658.
29. Fride E. Multiple roles for the endocannabinoid system during the earliest stages of life: pre- and postnatal development. *J Neuroendocrinol* 2008;20(suppl.1):75-81.
30. Watkins BA. Diet, endocannabinoids, and health. *Nutr Res* 2019 jun 15 [Epub ahead of print].
31. Merrit TA, Wilkinson B, Chervenak C. Maternal use of marijuana during pregnancy and lactation: implications for infant and child development and their well-being. *Acad J Ped Neonatol* 2016;2:1-7.
32. Alshaarawy O, Anthony JC. Cannabis use among women of reproductive age in the United States: 2002-2017. *Addict Behav* 2019;99:1-7.
33. Jaques SC, Kingsbury A, Henshcke P, Chomchai C, Clews S, Falconer J et al. Cannabis, the pregnant woman and her child: weedin out de myths. *J Perinatol* 2014;34:417-424.
34. Metz TD, Borgelt LM. Marijuana use in pregnancy and while breastfeeding. *Obstet Gynecol* 2018;132:1198-1212.
35. Foster BC, Abramovici H, Harris CS. Cannabis and cannabinoids: kinetics and interactions. *Am J Med* 2019. [Epub ahead of print].
36. Seltenrich N. Cannabis contaminants: regulating solvents, microbes, and metals in legal weed. *Environ Health Perspect* 2019;127:1-6.
37. Craven CB, Wawryk N, Jiang P, Liu Z, Li XF. Pesticides and trace elements in cannabis: analytical and environmental challenges and opportunities. *J Environ Sci* 2019;85:82-93.
38. Dryburgh LM, Bolan NS, Grof CPL, Galettis P, Schneider J, Lucas CJ et al. Cannabis contaminants: sources, distribution, human toxicity and pharmacologic effects. *Br J Clin Pharmacol* 2018;84:2468-2476.
39. Drummer OH, Gerastamoulos D, Woodford NW. Cannabis as a cause of death: a review. *Forensic Sci Int* 2019;298:298-306.
40. Budney AJ, Borodovsky JT. The potential impact of cannabis legalization on the development of cannabis use disorders. *Prev Med* 2017;104:31-36.
41. Richardson KA, Hester AK, McLemore GL. Prenatal cannabis exposure – the “first hit” to the endocannabinoid system. *Neurotoxicol Teratol* 2016;58:5-14.
42. Costa MA. The endocannabinoid system: a novel player in human placentation. *Reprod Toxicol* 2016;61:58-67.
43. Stickrath E. Marijuana use in pregnancy: an updated look at marijuana use and its impact on pregnancy. *Clin Obstet Gynecol* 2019;62:185-190.
44. Pressman P, Clemens RA. Neurodevelopment and exposure to cannabis. *Nutr Today* 2019;54:116-121.
45. Foeller ME, Lyell DJ. Marijuana use in pregnancy: concerns in an evolving era. *J Midw Womens Health* 2017;62:363-367.
46. Silverstein M, Howell EA, Zuckerman B. Cannabis use in pregnancy: a table of 2 concerns. *JAMA* 2019;322:121-122.
47. Calvigioni D, Hurd HL, Harkany T, Keimpema E. Neuronal substrates and functional consequences of prenatal cannabis exposure. *Eur Child Adolesc Psychiatry* 2014;23:931-941.
48. Metz TD, Stickrath EH. Marijuana use in pregnancy and lactation: a review of the evidence. *Am J Obstet Gynecol* 2015;213:761-778.
49. Warshak CR, Regan J, Moore B, Magner K, Kritzer S, Van Hook J. Association between marijuana use and adverse obstetrical and neonatal outcomes. *J Perinatol* 2015;35:991-995.
50. Lamy S, Laqueille X, Thibaut F. Conséquences potentielles de la consommation de tabac, de cannabis et de cocaïne par la femme enceinte sur la grossesse, le nouveau-né et l'enfant: revue de littérature. *L'Encephale* 2015;41:S13-S20.
51. Luke S, Hutcheon J, Kendall T. Cannabis use in pregnancy in British Columbia and selected birth outcomes. *J Obstet Gynaecol Can* 2019 Feb 7. pii: S1701-2163(18)30909-5. [Epub ahead of print].
52. Metz TD. What is new in cannabis use in pregnancy? *Obstet Gynecol* 2018;131:594-595.
53. Neradugomma NK, Drafton K, O'Day DR, Liao MZ, Han LW, Glass IA et al. Marijuana use differentially affects cannabinoid receptor expression in early gestational human endometrium and placenta. *Placenta* 2018;66:36-39.
54. Friedrich J, Khatib D, Parsa K, Santopietro A, Gallicano GI. The grass isn't always greener: The effects of cannabis on embryological development. *BMC Pharmacol Toxicol* 2016;17:45-57.
55. Youg-Wolf KC, Savovar V, Tucker L, Conway A, Alexeef S, Weisner C et al. Self-reported daily, weekly, and monthly cannabis use among women before and during pregnancy. *JAMA* 2019;2:1-10.
56. Ford TC, Hayley AC, Downey LA, Parrott AC. Cannabis: An overview of its adverse acute and chronic effects and its implications. *Curr Drug Abuse Rev* 2017;10:6-18.
57. Corsi DJ, Walsh L, Weiss D, Hsu H, El-Chaar D, Hawken S et al. Association between self-reported prenatal cannabis use and maternal, perinatal, and neonatal outcomes. *JAMA* 2019;322:145-152.
58. Schreiber S, Pick CG. Cannabis use during pregnancy: are we at the verge of defining a “fetal cannabis spectrum disorder”? *Med Hypotheses* 2019;124:53-55.

59. Jansson LM, Jordan CJ, Velez ML. Perinatal marijuana use and the developing child. *JAMA* 2018;320:545-546.
60. Leyton M. Cannabis legalization: did we make a mistake? Update 2019. *J Psychiatry Neurosci* 2019;44:291-293.
61. Crume TL, Juhl AL, Brooks-Russel A, Hall KE, Wymore E, Borgelt LM. Cannabis use during perinatal period in a state with legalized recreational and medical marijuana: the association between maternal characteristics, breastfeeding patterns, and neonatal outcomes. *J Pediatr* 2018;197:90-96.
62. Chang JC, Tarr JA, Holland CL, De Genna NM, Richardson GA, Rodriguez KL et al. Beliefs and attitudes regarding prenatal marijuana use: perspectives of pregnant women who report use. *Drug Alcohol Dep* 2019;196:14-20.
63. Chawla D et al. An epidemiological, developmental and clinical overview of cannabis use during pregnancy. *Prev Med* 2018;116:1-5.
64. AWHONN. Marijuana use during pregnancy. *J Obstet Gynecol Neonat Nurs* 2018;47:719-721.
65. Warner TD, Roussos-Ross D, Behnke M. Effects on maternal-fetal health and the developing child. *Clin Perinatol* 2014;41:877-894.